



75 Victor Heights Parkway STE C
Victor, NY 14564
Ph: 585.337.4300
Fax: 585.396.7264
www.medicineshoppepharm.com
www.compoundingrochester.com

Electronic Prescribing (E-Prescribing) Compounded Injectable Semaglutide

1. Identify our Pharmacy in your EMR system.
 - a. The Medicine Shoppe in Victor, NY
 - b. CRM-Canandaigua LLC in Victor, NY (some systems use our corporate name)
 - c. (585) 337-4300
 - d. 75 Victor Heights Parkway, STE C, Victor, NY 14564
 - e. Our NPI: 1427315266
2. Select the Medication
 - a. If your EMR system allows you to free-form a compounded medication that does not have a single-number NDC, consider free-form typing as the prescribed medication “**Compounded Semaglutide 2.5mg/ml**”
 - b. If your EMR will now allow you to free form, call the pharmacy and we can assist you with a verbal or fax Rx.
3. Add the Directions or “SIG” (used to embed compound name and directions – PICK ONE)
 - a. **Titration Dosage of Compounded injectable Semaglutide 2.5mg/ml**: Once weekly, inject 0.1ml for 4 weeks, then inject once weekly 0.2ml for 4 weeks. Quantity to be dispensed, 1ml.
 - b. **Ongoing Dosage of Compounded injectable Semaglutide 2.5mg/ml**: *(for 1mg / week)* Once weekly, inject 0.4ml. Quantity to be dispensed, 2ml.
4. Enter the authorized # of refills.
5. We also need an **Rx for Syringes to administer the Semaglutide Compound.**
 - a. Select a “27G x 1/2 ML TB Syringe”; Quantity to be dispensed: #10; Enter # of Refills.
 - b. Sig: To be used with Semaglutide Injection Weekly

**Call any of our pharmacists Mon-Fri from 9am – 5pm if you need assistance.
(585) 337-4300 Option #3, then #2.**



75 Victor Heights Parkway STE C
Victor, NY 14564
Ph: 585.337.4300
Fax: 585.396.7264
www.medicineshoppepharm.com
www.compoundingrochester.com

This is based upon published dosing for other branded Semaglutide GLP-1 medications. This chart indicates the sizes of the medication that we will dispense indicated by total volume (ml).

As always, these are recommendations, and you are free to dose as you choose. If you have further questions, please all and speak to any of our pharmacists.

Recommended Dosing	Dose (mg)	Volume Syringe (ml)	Total Volume (ml)
Mo #1 Weeks 1 - 4 0.25 mg / week	0.25	0.1	1
Mo #2 Weeks 5 - 8 0.5 mg / week	0.50	0.2	1
Mo #3 Weeks 9-12 1mg / week	1.00	0.4	2
Ongoing 1.7mg / week	1.70	0.68	4
Ongoing 2.4mg / week	2.40	0.96	4

- Weeks 1-7 can be completed with a 1ml vial.
- Week #8 - #12 will be filled with the 2ml vial for ongoing doses.
- If writing for 1.7mg/week or 2.4mg/week, we will dispense a 4ml vial which will last the patient 4-5 weeks.

No compounded Medication is reviewed by FDA for safety or efficacy. Above is general proposed guidance that should be adapted to each individual prescriber's practice.